

### **The influence of incident management teams on the deployment of wildfire suppression resources**

[www.nrfirescience.org/resource/15494](http://www.nrfirescience.org/resource/15494)

Despite large commitments of personnel and equipment to wildfire suppression, relatively little is known about the factors that affect how many resources are ordered and assigned to wildfire incidents and the variation in resources across incident management teams (IMTs). Using detailed data on suppression resource assignments for...

Year Published: 2017

Type: Document

Book or Chapter or Journal Article

### **Characterising resource use and potential inefficiencies during large-fire suppression in the western US**

[www.nrfirescience.org/resource/15492](http://www.nrfirescience.org/resource/15492)

Currently, limited research on large-fire suppression effectiveness suggests fire managers may over-allocate resources relative to values to be protected. Coupled with observations that weather may be more important than resource abundance to achieve control objectives, resource use may be driven more by risk aversion than...

Author(s): Hari Katuwal, Christopher J. Dunn, David E. Calkin

Year Published: 2017

Type: Document

Book or Chapter or Journal Article

### **Impacts of salvage logging on biodiversity – A meta-analysis**

[www.nrfirescience.org/resource/15274](http://www.nrfirescience.org/resource/15274)

Logging to 'salvage' economic returns from forests impacted by natural disturbances has become increasingly prevalent globally. Despite potential negative effects on biodiversity, salvage logging is often conducted, even in areas otherwise excluded from logging and reserved for nature conservation, inter alia because...

Author(s): Simon Thorn, Claus Bassler, Roland Brandl, Philip J. Burton, John L. Campbell, Rebecca Cahall, Jorge Castro, Chang-Yong Choi, Tyler Cobb, Daniel C. Donato, Ewa Durska, Joseph B. Fontaine, Sylvie Gauthier, Christian Hebert, Torsten Hothorn, Richard L. Hutto, Eun-Jae Lee, Alexandro B. Leverkus, David B. Lindenmayer, Martin K. Obrist, Josep Rost, Sebastian Seibold, Rupert Seidl, Dominik Thom, Kaysandra Waldron, Beat Wermelinger, Maria-Barbara Winter, Michal Zmihorski, Jorg Muller

Year Published: 2017

Type: Document

Book or Chapter or Journal Article

### **The influence of incident management teams on the deployment of wildfire suppression resources**

[www.nrfirescience.org/resource/15493](http://www.nrfirescience.org/resource/15493)

Despite large commitments of personnel and equipment to wildfire suppression, relatively little is known about the factors that affect how many resources are ordered and assigned to wildfire incidents and the variation in resources across incident management teams (IMTs). Using detailed data on suppression resource assignments for...

Author(s): Michael S. Hand, Hari Katuwal, David E. Calkin, Matthew P. Thompson

Year Published: 2017

Type: Document

Book or Chapter or Journal Article

### **Mixed-severity fire and salvage logging in dry forests of Oregon's western Cascades**

[www.nrfirescience.org/resource/15054](http://www.nrfirescience.org/resource/15054)

Interest in PNW forests is shifting from a focus on old-growth forests alone to include the ecological value and processes of early-seral communities. However, focusing on the alpha and omega states of a linear successional model does not account for the suite of conditions derived from mixed-severity fire common in many forests....

Author(s): Christopher J. Dunn, John D. Bailey

Year Published: 2017

Type: Document

Technical Report or White Paper

### **Effects of post-fire logging on fuel dynamics in a mixed-conifer forest, Oregon, USA: a 10-year assessment**

[www.nrfirescience.org/resource/14429](http://www.nrfirescience.org/resource/14429)

Removal of fire-killed trees (i.e. post-fire or salvage logging) is often conducted in part to reduce woody fuel loads and mitigate potential reburn effects. Studies of post-salvage fuel dynamics have primarily used chronosequence or modelling approaches, with associated limitations; longitudinal studies tracking fuels over time...

Author(s): John L. Campbell, Daniel C. Donato, Joseph B. Fontaine

Year Published: 2016

Type: Document

Book or Chapter or Journal Article

### **Response of native versus exotic plant guilds to cattle and elk herbivory in forested rangeland**

[www.nrfirescience.org/resource/13843](http://www.nrfirescience.org/resource/13843)

Are exotic plant species favoured by non-native ungulate herbivores and disadvantaged by native herbivores in forested rangelands? Do the impacts of ungulates on exotic vs native plants depend on forest management activities such as prescribed fire and stand thinning? Location: Northeastern Oregon, USA. Methods: We recorded changes...

Author(s): Burak K. Pekin, Michael J. Wisdom, Catherine G. Parks, Bryan A. Endress, Bridgett J. Naylor

Year Published: 2016

Type: Document

Book or Chapter or Journal Article

### **The integrated rangeland fire management strategy actionable science plan**

[www.nrfirescience.org/resource/14697](http://www.nrfirescience.org/resource/14697)

The Integrated Rangeland Fire Management Strategy (hereafter Strategy, DOI 2015) outlined the need for coordinated, science-based adaptive management to achieve long-term protection, conservation, and restoration of the sagebrush (*Artemisia* spp.) ecosystem. A key component of this management approach is the...

Author(s): Integrated Rangeland Fire Management Strategy Actionable Science Plan Team

Year Published: 2016

Type: Document

Management or Planning Document

### **Post-fire logging produces minimal persistent impacts on understory vegetation in northeastern Oregon, USA**

[www.nrfirescience.org/resource/14354](http://www.nrfirescience.org/resource/14354)

Post-fire forest management commonly requires accepting some negative ecological impacts from management activities in order to achieve management objectives. Managers need to know, however, whether ecological impacts from post-fire management activities are transient or cause long-term

ecosystem degradation. We studied the long-...  
Author(s): David W. Peterson, Erich K. Dodson  
Year Published: 2016  
Type: Document  
Book or Chapter or Journal Article

**Rapid response tools and datasets for post-fire modeling: linking earth observations and process-based hydrological models to support post-fire remediation**

[www.nrfirescience.org/resource/15538](http://www.nrfirescience.org/resource/15538)

Post-wildfire flooding and erosion can threaten lives, property and natural resources. Increased peak flows and sediment delivery due to the loss of surface vegetation cover and fire-induced changes in soil properties are of great concern to public safety. Burn severity maps derived from remote sensing data reflect fire-induced...

Author(s): Mary Ellen Miller, Michael Billmire, William J. Elliot, Kevin A. Endsley, Peter R. Robichaud  
Year Published: 2016  
Type: Document  
Book or Chapter or Journal Article

**Rapid-response tools and datasets for post-fire remediation: linking remote sensing and process-based hydrological models**

[www.nrfirescience.org/resource/14641](http://www.nrfirescience.org/resource/14641)

Post-wildfire flooding and erosion can threaten lives, property and natural resources. Increased peak flows and sediment delivery due to the loss of surface vegetation cover and fire-induced changes in soil properties are of great concern to public safety. Burn severity maps derived from remote sensing data reflect fire-induced...

Author(s): Mary Ellen Miller, William J. Elliot, Peter R. Robichaud, Kevin A. Endsley  
Year Published: 2016  
Type: Document  
Book or Chapter or Journal Article

**Ecological Effects of Post-fire Salvage Logging in the Pacific Northwest**

[www.nrfirescience.org/resource/15729](http://www.nrfirescience.org/resource/15729)

Post-fire salvage logging is typically proposed as a means of recovering some of the lost economic value in dead or damaged trees. The ecological consequences of salvage, however, are often considered negative from the perspective of soils, hydrology, and wildlife habitat resources, although species responses do vary. Early...

Author(s): Matthew J. Reilly, Thomas A. Spies, Paul F. Hessburg  
Year Published: 2015  
Type: Document  
Research Brief or Fact Sheet

**Rapid response tools and datasets for post-fire modeling: linking earth observations and process-based hydrological models to support post-fire remediation**

[www.nrfirescience.org/resource/13466](http://www.nrfirescience.org/resource/13466)

Preparation is key to utilizing Earth Observations and process-based models to support post-wildfire mitigation. Post-fire flooding and erosion can pose a serious threat to life, property and municipal water supplies. Increased runoff and sediment delivery due to the loss of surface cover and fire-induced changes in soil...

Author(s): Mary Ellen Miller, Michael Billmire, William J. Elliot, Kevin A. Endsley, Peter R. Robichaud  
Year Published: 2015  
Type: Document

**Assessing soil and vegetation recovery following the 2005 School Fire, Umatilla National Forest - 10-year update**

[www.nrfirescience.org/resource/12811](http://www.nrfirescience.org/resource/12811)

Following the 2005 School Fire which burned ~ 50,000 acres across forest and grasslands, managers were particularly concerned with treating severely burned areas to mitigate weed spread and to limit soil erosion. Various mulching treatments (wheat straw, wood strand, and hydromulch) were implemented to control...

Author(s): Peter R. Robichaud, Penelope Morgan, Leigh B. Lentile, Sarah A. Lewis, Andrew T. Hudak, Deborah S. Page-Dumroese

Year Published: 2015

Type: Document

Research Brief or Fact Sheet

**Vegetation response to burn severity, native grass seeding, and salvage logging**

[www.nrfirescience.org/resource/13422](http://www.nrfirescience.org/resource/13422)

As the size and extent of wildfires has increased in recent decades, so has the cost and extent of post-fire management, including seeding and salvage logging. However, we know little about how burn severity, salvage logging, and post-fire seeding interact to influence vegetation recovery long-term. We sampled understory plant...

Author(s): Penelope Morgan, Marshall Moy, Christine A. Droske, Leigh B. Lentile, Sarah A. Lewis, Peter R. Robichaud, Andrew T. Hudak, Christopher Jason Williams

Year Published: 2015

Type: Document

Book or Chapter or Journal Article

**Effects of post-fire salvage logging and a skid trail treatment on ground cover, soils, and sediment production in the interior western United States**

[www.nrfirescience.org/resource/12829](http://www.nrfirescience.org/resource/12829)

Post-fire salvage logging adds another set of environmental effects to recently burned areas, and previous studies have reported varying impacts on vegetation, soil disturbance, and sediment production with limited data on the underlying processes. Our objectives were to determine how: (1) ground-based post-fire logging affects...

Author(s): Joseph W. Wagenbrenner, Lee H. MacDonald, Robert N. Coats, Peter R. Robichaud, Robert E. Brown

Year Published: 2015

Type: Document

Book or Chapter or Journal Article

**Prescribed fire effects on resource selection by cattle in mesic sagebrush steppe. part 1: spring grazing**

[www.nrfirescience.org/resource/12148](http://www.nrfirescience.org/resource/12148)

Prescribed fire is commonly applied world-wide as a tool for enhancing habitats and altering resource-selection patterns of grazing animals. A scientific basis for this practice has been established in some ecosystems but its efficacy has not been rigorously evaluated on mesic sagebrush steppe. Beginning in 2003, resource-selection...

Author(s): Patrick E. Clark, Jaechoul Lee, Kyungduk Ko, Ryan M. Nielson, Douglas E. Johnson, David C. Ganskopp, Joe Chigbrow, Frederick B. Pierson, Stuart P. Hardegree

Year Published: 2014

Type: Document

Book or Chapter or Journal Article

### **Is proportion burned severely related to daily area burned?**

[www.nrfirescience.org/resource/13018](http://www.nrfirescience.org/resource/13018)

The ecological effects of forest fires burning with high severity are long-lived and have the greatest impact on vegetation successional trajectories, as compared to low-to-moderate severity fires. The primary drivers of high severity fire are unclear, but it has been hypothesized that wind-driven, large fire-growth days play a...

Author(s): Donovan Birch, Penelope Morgan, Crystal A. Kolden, Andrew T. Hudak, Alistair M. S. Smith

Year Published: 2014

Type: Document

Book or Chapter or Journal Article

### **Using resistance and resilience concepts to reduce impacts of invasive annual grasses and altered fire regimes on the sagebrush ecosystem and greater sage-grouse: a strategic multi-scale approach**

[www.nrfirescience.org/resource/12989](http://www.nrfirescience.org/resource/12989)

This Report provides a strategic approach for conservation of sagebrush ecosystems and Greater Sage- Grouse (sage-grouse) that focuses specifically on habitat threats caused by invasive annual grasses and altered fire regimes. It uses information on factors that influence (1) sagebrush ecosystem resilience to disturbance and...

Author(s): Jeanne C. Chambers, David A. Pyke, Jeremy D. Maestas, Michael L. Pellant, Chad S. Boyd, Steven B. Campbell, Shawn Espinosa, Douglas W. Havlina, Kenneth E. Mayer, Amarina Wuenschel

Year Published: 2014

Type: Document

Management or Planning Document

### **A synthesis of post-fire Burned Area Reports from 1972 to 2009 for western US Forest Service lands: trends in wildfire characteristics and post-fire stabilisation treatments and expenditures**

[www.nrfirescience.org/resource/13010](http://www.nrfirescience.org/resource/13010)

Over 1200 post-fire assessment and treatment implementation reports from four decades (1970s-2000s) of western US forest fires have been examined to identify decadal patterns in fire characteristics and the justifications and expenditures for the post-fire treatments. The main trends found were: (1) the area burned by wildfire...

Author(s): Peter R. Robichaud, Hakjun Rhee, Sarah A. Lewis

Year Published: 2014

Type: Document

Book or Chapter or Journal Article, Synthesis

### **Semiarid rangeland is resilient to summer fire and postfire grazing utilization**

[www.nrfirescience.org/resource/12050](http://www.nrfirescience.org/resource/12050)

Most wildfires occur during summer in the northern hemisphere, the area burned annually is increasing, and fire effects during this season are least understood. Understanding plant response to grazing following summer fire is required to reduce ecological and financial risks associated with wildfire. Forty 0.75-ha plots were...

Author(s): Lance T. Vermeire, Jessica L. Crowder, David B. Wester

Year Published: 2014

Type: Document

Book or Chapter or Journal Article

### **Quantifying restoration effectiveness using multi-scale habitat models: implications for sage-grouse in the Great Basin**

[www.nrfirescience.org/resource/12947](http://www.nrfirescience.org/resource/12947)

A recurrent challenge in the conservation of wide-ranging, imperiled species is understanding which habitats to protect and whether we are capable of restoring degraded landscapes. For Greater Sage-grouse (*Centrocercus urophasianus*), a species of conservation concern in the western United States, we approached this problem by...

Author(s): Robert S. Arkle, David S. Pilliod, Steven E. Hanser, Matthew L. Brooks, Jeanne C. Chambers, James B. Grace, Kevin C. Knutson, David A. Pyke, Justin L. Welty, Troy A. Wirth

Year Published: 2014

Type: Document

Book or Chapter or Journal Article

### **The Bitterroot Valley fires of 2000 - Revisiting experiences and fire effects 13 years later**

[www.nrfirescience.org/resource/12673](http://www.nrfirescience.org/resource/12673)

During the Fires of 2000 field trip, held as part of the May 2014 Large Wildland Fires Conference, researchers, managers, residents, and stakeholders shared their experiences around the unprecedented number and size of fires that burned in the Bitterroot Valley in the summer of 2000. Topics discussed included fire history, fire...

Author(s): Corey L. Gucker

Year Published: 2014

Type: Document

Research Brief or Fact Sheet

### **Using native annual plants to restore post-fire habitats in western North America**

[www.nrfirescience.org/resource/12139](http://www.nrfirescience.org/resource/12139)

Increasing fire frequencies and uncharacteristic severe fires have created a need for improved restoration methods across rangelands in western North America. Traditional restoration seed mixtures of native perennial mid- to late-seral plant species may not be suitable for intensely burned sites that have been returned to an early-...

Author(s): Christopher M. Herron, Jayne L. Jonas, Paul J. Meiman, Mark W. Paschke

Year Published: 2013

Type: Document

Book or Chapter or Journal Article

### **Perspectives on disconnects between scientific information and management decisions on post-fire recovery in western US**

[www.nrfirescience.org/resource/12035](http://www.nrfirescience.org/resource/12035)

Environmental regulations frequently mandate the use of 'best available' science, but ensuring that it is used in decisions around the use and protection of natural resources is often challenging. In the Western US, this relationship between science and management is at the forefront of post-fire land management decisions. Recent...

Author(s): Xiaoli Chen, Nathan Emery, Elizabeth S. Garcia, Erin J. Hanan, Heather E. Hodges, Tyrone Martin, Matthew A. Meyers, Lindsey E. Peavey, Hui Peng, Jaime Sainz Santamaria, Kellie A. Uyeda, Sarah E. Anderson, Christina Tague

Year Published: 2013

Type: Document

Book or Chapter or Journal Article

### **Post-fire mulching for runoff and erosion mitigation; Part I: effectiveness at reducing hillslope erosion rates**



[www.nrfirescience.org/resource/11994](http://www.nrfirescience.org/resource/11994)

Mulch treatments often are used to mitigate post-fire increases in runoff and erosion rates but the comparative effectiveness of various mulches is not well established. The ability of mulch treatments to reduce sediment yields from natural rainfall and resulting overland flow was measured using hillslope plots on areas burned at...

Author(s): Peter R. Robichaud, Sarah A. Lewis, Joseph W. Wagenbrenner, Louise E. Ashmun, Robert E. Brown

Year Published: 2013

Type: Document

Book or Chapter or Journal Article

### **Effectiveness of post-fire Burned Area Emergency Response (BAER) road treatments: results from three wildfires**

[www.nrfirescience.org/resource/12142](http://www.nrfirescience.org/resource/12142)

Wildland fires often cause extreme changes in the landscape that drastically influence surface runoff and soil erosion, which can impact forest resources, aquatic habitats, water supplies, public safety, and forest access infrastructure such as forest roads. Little information is available on the effectiveness of various post-fire...

Author(s): Randy B. Foltz, Peter R. Robichaud

Year Published: 2013

Type: Document

Technical Report or White Paper

### **Does seeding after wildfires in rangelands reduce erosion or invasive species?**

[www.nrfirescience.org/resource/12132](http://www.nrfirescience.org/resource/12132)

Mitigation of ecological damage caused by rangeland wildfires has historically been an issue restricted to the western United States. It has focused on conservation of ecosystem function through reducing soil erosion and spread of invasive plants. Effectiveness of mitigation treatments has been debated recently. We reviewed recent...

Author(s): David A. Pyke

Year Published: 2013

Type: Document

Book or Chapter or Journal Article, Synthesis

### **Climate change, forests, fire, water, and fish: building resilient landscapes, streams, and managers**

[www.nrfirescience.org/resource/11270](http://www.nrfirescience.org/resource/11270)

Fire will play an important role in shaping forest and stream ecosystems as the climate changes. Historic observations show increased dryness accompanying more widespread fire and forest die-off. These events punctuate gradual changes to ecosystems and sometimes generate stepwise changes in ecosystems. Climate vulnerability...

Author(s): Charles H. Luce, Penelope Morgan, Kathleen A. Dwire, Daniel J. Isaak, Zachary A. Holden, Bruce E. Rieman

Year Published: 2012

Type: Document

Synthesis, Technical Report or White Paper

### **Wildland fire in ecosystems: effects of fire on cultural resources and archaeology**

[www.nrfirescience.org/resource/141](http://www.nrfirescience.org/resource/141)

This state-of-knowledge review provides a synthesis of the effects of fire on cultural resources, which can be used by fire managers, cultural resource (CR) specialists, and archaeologists to more effectively

manage wildland vegetation, fuels, and fire. The goal of the volume is twofold: (1) to provide cultural resource/...

Year Published: 2012

Type: Document

Technical Report or White Paper

### **Assessing the success of postfire reseeding in semiarid rangelands using terra MODIS**

[www.nrfirescience.org/resource/11489](http://www.nrfirescience.org/resource/11489)

Successful post-fire reseeding efforts may aid rangeland ecosystem recovery by rapidly establishing a desired plant community and thereby reducing the likelihood of infestation by invasive plants. While the success of post-fire remediation is critical, few efforts have been made to leverage existing geospatial technologies to...

Author(s): Fang Chen, Keith T. Weber, John L. Schnase

Year Published: 2012

Type: Document

Book or Chapter or Journal Article

### **Utility of remotely sensed imagery for assessing the impact of salvage logging after forest fires**

[www.nrfirescience.org/resource/8352](http://www.nrfirescience.org/resource/8352)

Remotely sensed imagery provides a useful tool for land managers to assess the extent and severity of post-wildfire salvage logging disturbance. This investigation uses high resolution QuickBird and National Agricultural Imagery Program (NAIP) imagery to map soil exposure after ground-based salvage operations. Three wildfires with...

Author(s): Sarah A. Lewis, Peter R. Robichaud, Andrew T. Hudak, Brian Austin, Robert J. Liebermann

Year Published: 2012

Type: Document

Book or Chapter or Journal Article

### **Using native annual plant species to suppress weedy invasive species in post-fire habitats - Final Report to the Joint Fire Science Program**

[www.nrfirescience.org/resource/11467](http://www.nrfirescience.org/resource/11467)

Increasing fire frequencies and uncharacteristic severe fires have created a need for improved restoration methods across rangelands in western North America. Traditional restoration seed mixtures of perennial mid- to late-seral plant species may not be suitable for intensely burned sites that have been returned to an early-seral...

Author(s): Mark W. Paschke, Paul J. Meiman, William H. Romme, Cynthia S. Brown

Year Published: 2011

Type: Document

Technical Report or White Paper

### **Effects of post-fire salvage logging on cavity-nesting birds and small mammals in southeastern Montana**

[www.nrfirescience.org/resource/12052](http://www.nrfirescience.org/resource/12052)

We investigated how post-fire salvage logging of Ponderosa Pine (*Pinus ponderosa*) affected populations of cavity-nesting birds and small mammals in southeastern Montana in 2004 and 2005. We examined two salvage and two control plots with three point-count stations and one small mammal trap site randomly distributed across each plot...

Author(s): William J. Kronland, Marco Restani

Year Published: 2011

Type: Document

Book or Chapter or Journal Article



### **Recent trends in post-wildfire seeding in western US forests: costs and seed mixes**

[www.nrfirescience.org/resource/8284](http://www.nrfirescience.org/resource/8284)

Broadcast seeding is one of the most commonly used post-fire rehabilitation treatments to establish ground cover for erosion control and mitigation of non-native plant species invasions. Little quantitative information is available on overall trends of post-fire seeding expenditures and seed mixes used over time in forested...

Author(s): Donna Peppin, Peter Z. Fule, Carolyn Hull Sieg, Jan L. Beyers, Molly E. Hunter, Peter R. Robichaud

Year Published: 2011

Type: Document

Book or Chapter or Journal Article, Synthesis

### **Does seeding after severe forest fires in western USA mitigate negative impacts on soils and plant communities?**

[www.nrfirescience.org/resource/11501](http://www.nrfirescience.org/resource/11501)

Broadcast seeding is one of the most widely used post-wildfire emergency response treatments intended to reduce soil erosion, increase vegetative ground cover, and minimize establishment and spread of non-native plant species. However, seeding treatments can also have negative effects such as competition with recovering native...

Author(s): Donna Peppin, Peter Z. Fule, Jan L. Beyers, Carolyn Hull Sieg, Molly E. Hunter

Year Published: 2011

Type: Document

Synthesis, Technical Report or White Paper

### **Woodpecker habitat after the fire**

[www.nrfirescience.org/resource/13508](http://www.nrfirescience.org/resource/13508)

Public land managers are asked to minimize fuel levels after fires, including using techniques such as salvage logging. They are also responsible for maintaining suitable wildlife habitat, especially for species of concern to state and federal agencies. An area where these responsibilities could conflict is in the use of salvage...

Author(s): Victoria A. Saab

Year Published: 2011

Type: Document

Research Brief or Fact Sheet

### **Getting results: measuring post-wildfire erosion control treatment effectiveness**

[www.nrfirescience.org/resource/11031](http://www.nrfirescience.org/resource/11031)

In the past decade, wildfires around the world have continued to increase in size, severity, and cost. The number of people living in wildland areas has also increased, putting public safety, homes, roads, public infrastructure, water quality, and valued natural resources at risk from wildfire and secondary fire effects. Major...

Author(s): Peter R. Robichaud, Robert E. Brown, Peter M. Wohlgenuth, Joseph W. Wagenbrenner

Year Published: 2011

Type: Document

Conference Proceedings

### **Greater sage-grouse: Ecology and conservation of a landscape species and its habitats**

[www.nrfirescience.org/resource/15406](http://www.nrfirescience.org/resource/15406)

The greater sage-grouse is at the center of a complex challenge to conserve sagebrush ecosystems.

The species has declined across much of its range, including 11 western states and 2 Canadian provinces, mostly due to loss of critical sagebrush habitat. Agriculture, roads, development of energy resources, wildfire, and invasive...

Author(s): Steve Knick, John W. Connelly

Year Published: 2011

Type: Document

Book or Chapter or Journal Article

### **Delaying sheep grazing after wildfire in sagebrush steppe may not affect vegetation recovery**

[www.nrfirescience.org/resource/11439](http://www.nrfirescience.org/resource/11439)

Although many land managers prohibit grazing for 2 years after a fire, little research has been conducted to determine the interaction of grazing with vegetation recovery after fire. In a study conducted in sagebrush steppe rangelands after a 2000 wildfire at the United States Sheep Experiment Station in Idaho, the influence of...

Author(s): Lovina Roselle, Steven S. Seefeldt, Karen Launchbaugh

Year Published: 2010

Type: Document

Book or Chapter or Journal Article

### **Post-fire treatment effectiveness for hillslope stabilization**

[www.nrfirescience.org/resource/12594](http://www.nrfirescience.org/resource/12594)

This synthesis of post-fire treatment effectiveness reviews the past decade of research, monitoring, and product development related to post-fire hillslope emergency stabilization treatments, including erosion barriers, mulching, chemical soil treatments, and combinations of these treatments. In the past ten years, erosion barrier...

Author(s): Peter R. Robichaud, Louise E. Ashmun, Bruce D. Sims

Year Published: 2010

Type: Document

Synthesis, Technical Report or White Paper

### **Continued evaluation of post-fire recovery and treatment effectiveness for validation of the ERMiT erosion model (combined proposals P07-2-2-10 and P07-2-3-06) - Final Report to the Joint Fire Science Program**

[www.nrfirescience.org/resource/11227](http://www.nrfirescience.org/resource/11227)

The use and cost of post-fire emergency stabilization treatments continues to grow. To help maximize the impact of these treatments, many assessment teams use the Erosion Risk Management Tool (ERMiT) erosion model to predict postfire erosion and mitigation effects. However, despite several completed JFSP projects, the long-term...

Author(s): Peter R. Robichaud, William J. Elliot, Joseph W. Wagenbrenner, Sarah A. Lewis, Louise E. Ashmun, Peter M. Wohlgenuth, Robert E. Brown

Year Published: 2010

Type: Document

Technical Report or White Paper

### **Post-wildfire seeding in forests of the western United States: an evidence-based review**

[www.nrfirescience.org/resource/12595](http://www.nrfirescience.org/resource/12595)

Broadcast seeding is one of the most widely used post-wildfire emergency response treatments intended to reduce soil erosion, increase vegetative ground cover, and minimize establishment and spread of non-native plant species. We conducted an evidence-based review to examine the effectiveness and effects of post-wildfire seeding...

Author(s): Donna Peppin, Peter Z. Fule, Carolyn Hull Sieg, Jan L. Beyers, Molly E. Hunter

Year Published: 2010  
Type: Document  
Book or Chapter or Journal Article, Synthesis

### **Emergency post-fire rehabilitation treatment effects on burned area ecology and long-term restoration**

[www.nrfirescience.org/resource/12591](http://www.nrfirescience.org/resource/12591)

The predicted continuation of strong drying and warming trends in the southwestern United States underlies the associated prediction of increased frequency, area, and severity of wildfires in the coming years. As a result, the management of wildfires and fire effects on public lands will continue to be a major land management...

Author(s): Peter R. Robichaud, Sarah A. Lewis, Robert E. Brown, Louise E. Ashmun

Year Published: 2009

Type: Document

Book or Chapter or Journal Article, Synthesis

### **Nest-site selection by cavity-nesting birds in relation to postfire salvage logging**

[www.nrfirescience.org/resource/8383](http://www.nrfirescience.org/resource/8383)

Large wildfire events in coniferous forests of the western United States are often followed by postfire timber harvest. The long-term impacts of postfire timber harvest on fire-associated cavity-nesting bird species are not well documented. We studied nest-site selection by cavity-nesting birds over a 10-year period (1994-2003),...

Author(s): Victoria A. Saab, Robin E. Russell, Jonathan G. Dudley

Year Published: 2009

Type: Document

Book or Chapter or Journal Article

### **A synthesis of postfire road treatments for BAER teams: methods, treatment effectiveness, and decisionmaking tools for rehabilitation**

[www.nrfirescience.org/resource/12622](http://www.nrfirescience.org/resource/12622)

We synthesized post-fire road treatment information to assist BAER specialists in making road rehabilitation decisions. We developed a questionnaire; conducted 30 interviews of BAER team engineers and hydrologists; acquired and analyzed gray literature and other relevant publications; and reviewed road rehabilitation procedures and...

Author(s): Randy B. Foltz, Peter R. Robichaud, Hakjun Rhee

Year Published: 2009

Type: Document

Synthesis, Technical Report or White Paper

### **Listening to the message of the Black-backed Woodpecker, a hot fire specialist**

[www.nrfirescience.org/resource/11083](http://www.nrfirescience.org/resource/11083)

The Black-backed Woodpecker is an uncommon bird of the northern coniferous forests of North America. It is one of several species of fauna that are considered fire specialists. This woodpecker nests in cavities it creates in dead standing trees and feeds on wood-boring beetles and their larvae, which are also attracted to stressed...

Author(s): Elise LeQuire

Year Published: 2009

Type: Document

Research Brief or Fact Sheet

### **Effects of timber harvest following wildfire in western North America**

[www.nrfirescience.org/resource/11122](http://www.nrfirescience.org/resource/11122)

Timber harvest following wildfire leads to different outcomes depending on the biophysical setting of the forest, pattern of burn severity, operational aspects of tree removal, and other management activities. Fire effects range from relatively minor, in which fire burns through the understory and may kill a few trees, to severe, in...

Author(s): David L. Peterson, James K. Agee, Gregory H. Aplet, Dennis P. Dykstra, Russell T. Graham, John F. Lehmkuhl, David S. Pilliod, Donald F. Potts, Robert F. Powers, John D. Stuart

Year Published: 2009

Type: Document

Technical Report or White Paper

### **Effectiveness of aerial seeding and straw mulch for reducing post-wildfire erosion, north-western Montana, USA**

[www.nrfirescience.org/resource/8200](http://www.nrfirescience.org/resource/8200)

Various methods are available to reduce post-wildfire erosion, but there is limited quantitative information on the relative effectiveness of these techniques. We used rainfall simulations to compare the erosion and runoff rates from adjacent 0.5-m<sup>2</sup> plots treated with aerial grass seeding and straw mulch with untreated control plots...

Author(s): Amy H. Groen, Scott W. Woods

Year Published: 2008

Type: Document

Book or Chapter or Journal Article

### **Wildland fire in ecosystems: fire and nonnative invasive plants**

[www.nrfirescience.org/resource/12531](http://www.nrfirescience.org/resource/12531)

This state-of-knowledge review of information on relationships between wildland fire and nonnative invasive plants can assist fire managers and other land managers concerned with prevention, detection, and eradication or control of nonnative invasive plants. The 16 chapters in this volume synthesize ecological and botanical...

Year Published: 2008

Type: Document

Synthesis, Technical Report or White Paper

### **Measuring effectiveness of three postfire hillslope erosion barrier treatments, western Montana, USA**

[www.nrfirescience.org/resource/8389](http://www.nrfirescience.org/resource/8389)

After the Valley Complex Fire burned 86 000 ha in western Montana in 2000, two studies were conducted to determine the effectiveness of contour-felled log, straw wattle, and hand-dug contour trench erosion barriers in mitigating postfire runoff and erosion. Sixteen plots were located across a steep, severely burned slope, with a...

Author(s): Peter R. Robichaud, Frederick B. Pierson, Robert E. Brown, Joseph W. Wagenbrenner

Year Published: 2008

Type: Document

Book or Chapter or Journal Article

### **Evaluating the effectiveness of contour-felled log erosion barriers as a post-fire runoff and erosion mitigation treatment in the western United States**

[www.nrfirescience.org/resource/8167](http://www.nrfirescience.org/resource/8167)

Between 1998 and 2002, six sites were established immediately after large wildfires in the western United States to determine the effectiveness of contour-felled log erosion barriers in mitigating post-

wildfire runoff and erosion. In each pair of matched, burned, and small watersheds (1-13 ha), one was treated with contour-felled...

Author(s): Peter R. Robichaud, Joseph W. Wagenbrenner, Robert E. Brown, Peter M. Wohlgemuth, Jan L. Beyers

Year Published: 2008

Type: Document

Book or Chapter or Journal Article

### **Nest densities of cavity-nesting birds in relation to postfire salvage logging and time since wildfire**

[www.nrfirescience.org/resource/8145](http://www.nrfirescience.org/resource/8145)

We monitored the nest densities and nest survival of seven cavity-nesting bird species, including four open-space foragers (American Kestrel [*Falco sparverius*], Lewis's Woodpecker [*Melanerpes lewis*], Western Bluebird [*Sialia mexicana*], and Mountain Bluebird [*S. currucoides*]) and three wood-foragers (Hairy Woodpecker [*Picoides*...

Author(s): Victoria A. Saab, Robin E. Russell, Jonathan G. Dudley

Year Published: 2007

Type: Document

Book or Chapter or Journal Article

### **Predicting postfire erosion and mitigation effectiveness with a web-based probabilistic erosion model**

[www.nrfirescience.org/resource/8138](http://www.nrfirescience.org/resource/8138)

The decision of where, when, and how to apply the most effective postfire erosion mitigation treatments requires land managers to assess the risk of damaging runoff and erosion events occurring after a fire. To meet this challenge, the Erosion Risk Management Tool (ERMiT) was developed. ERMiT is a web-based application that uses the...

Author(s): Peter R. Robichaud, William J. Elliot, Frederick B. Pierson, David E. Hall, Corey A. Moffet

Year Published: 2007

Type: Document

Book or Chapter or Journal Article

### **Snag longevity in relation to wildfire and postfire salvage logging**

[www.nrfirescience.org/resource/8142](http://www.nrfirescience.org/resource/8142)

Snags create nesting, foraging, and roosting habitat for a variety of wildlife species. Removal of snags through postfire salvage logging reduces the densities and size classes of snags remaining after wildfire. We determined important variables associated with annual persistence rates (the probability a snag remains standing from 1...

Author(s): Robin E. Russell, Victoria A. Saab, Jonathan G. Dudley, Jay J. Rotella

Year Published: 2006

Type: Document

Book or Chapter or Journal Article

### **Protection from erosion following wildfire**

[www.nrfirescience.org/resource/11053](http://www.nrfirescience.org/resource/11053)

Erosion in the first year after a wildfire can be up to three orders of magnitude greater than the erosion from undisturbed forests. To mitigate potential postfire erosion, various erosion control treatments are applied on highly erodible areas with downstream resources in need of protection. Because postfire erosion rates generally...

Author(s): Peter R. Robichaud, William J. Elliot

Year Published: 2006

Type: Document  
Conference Proceedings

### **Postfire logging in riparian ecosystems**

[www.nrfirescience.org/resource/8126](http://www.nrfirescience.org/resource/8126)

We reviewed the behavior of wildfire in riparian zones, primarily in the western United States, and the potential ecological consequences of postfire logging. Fire behavior in riparian zones is complex, but many aquatic and riparian organisms exhibit a suite of adaptations that allow relatively rapid recovery after fire. Unless...

Author(s): Gordon H. Reeves, Peter A. Bisson, Bruce E. Rieman, Lee E. Benda

Year Published: 2006

Type: Document

Book or Chapter or Journal Article

### **Monitoring changes in soil quality from post-fire logging in the inland northwest**

[www.nrfirescience.org/resource/11015](http://www.nrfirescience.org/resource/11015)

The wildland fires of 2000, 2002, and 2003 created many opportunities to conduct post-fire logging operations in the Inland Northwest. Relatively little information is available on the impact of post-fire logging on long-term soil productivity or on the best method for monitoring these changes. We present a USDA Forest Service...

Author(s): Deborah S. Page-Dumroese, Martin F. Jurgensen, Ann Abbott, Thomas M. Rice, Joanne M. Tirocke, Sue Farley, Sharon DeHart

Year Published: 2006

Type: Document

Conference Proceedings

### **The effects of postfire salvage logging on cavity-nesting birds**

[www.nrfirescience.org/resource/12933](http://www.nrfirescience.org/resource/12933)

We investigated the effects of postfire salvage logging on cavity-nesting birds by comparing nest densities and patterns of nest reuse over a three-year period in seven logged and eight unlogged patches of mixed-conifer forest in the Blackfoot-Clearwater Wildlife Management Area, Montana. We found 563 active nests of 18 cavity-...

Author(s): Richard L. Hutto, Susan M. Gallo

Year Published: 2006

Type: Document

Book or Chapter or Journal Article

### **Post-fire logging debate ignores many issues**

[www.nrfirescience.org/resource/14599](http://www.nrfirescience.org/resource/14599)

Recent controversy concerning post-fire logging in Oregon is emblematic of the problems of "salvage logging" globally. Although tree regeneration after disturbances in forested areas is important, a narrow view of this issue ignores important ecological lessons, especially the role of disturbances in diversifying and rejuvenating...

Author(s): Dominick A. DellaSala, James R. Karr, Tania L. Schoennagel, David A. Perry, Reed F. Noss, David B. Lindenmayer, Robert L. Beschta, Richard L. Hutto, Mark E. Swanson, Jon Evans

Year Published: 2006

Type: Document

Book or Chapter or Journal Article

### **Toward meaningful snag-management guidelines for postfire salvage logging in North American**



### **conifer forests**

[www.nrfirescience.org/resource/14507](http://www.nrfirescience.org/resource/14507)

The bird species in western North America that are most restricted to, and therefore most dependent on, severely burned conifer forests during the first years following a fire event depend heavily on the abundant standing snags for perch sites, nest sites, and food resources. Thus, it is critical to develop and apply appropriate...

Author(s): Richard L. Hutto

Year Published: 2006

Type: Document

Book or Chapter or Journal Article

### **Wildfire, timber salvage, and the economics of expediency**

[www.nrfirescience.org/resource/8124](http://www.nrfirescience.org/resource/8124)

Administrative planning rules and legal challenges can have significant economic impacts on timber salvage programs on public lands. This paper examines the costs of the delay in salvage caused by planning rules and the costs associated with the volume reductions forced by legal challenges in one case study. The fires on the...

Author(s): Jeffrey P. Prestemon, David N. Wear, Fred J. Stewart, Thomas P. Holmes

Year Published: 2006

Type: Document

Book or Chapter or Journal Article

### **Fire management impacts on invasive plants in the western United States**

[www.nrfirescience.org/resource/12024](http://www.nrfirescience.org/resource/12024)

Fire management practices affect alien plant invasions in diverse ways. I considered the impact of six fire management practices on alien invasions: fire suppression, forest fuel reduction, prescription burning in crown-fire ecosystems, fuel breaks, targeting of noxious aliens, and postfire rehabilitation. Most western United States...

Author(s): Jon E. Keeley

Year Published: 2006

Type: Document

Book or Chapter or Journal Article, Synthesis

### **Assessing the causes, consequences and spatial variability of burn severity: a rapid response proposal - Final Report to the Joint Fire Science Program**

[www.nrfirescience.org/resource/11149](http://www.nrfirescience.org/resource/11149)

In this rapid response project, we have collected data on post-fire effects and pre-fire fuels and vegetation from 10 large fires that burned in 2003 and 2004. We use field and remotely sensed data collected during and soon after wildfires to quantify the interactions and spatial variability in fire effects, fuels, fire behavior,...

Author(s): Penelope Morgan, Andrew T. Hudak, Peter R. Robichaud, Kevin C. Ryan

Year Published: 2005

Type: Document

Technical Report or White Paper

### **Home range size and foraging habitat of Black-backed Woodpeckers**

[www.nrfirescience.org/resource/11417](http://www.nrfirescience.org/resource/11417)

We examined home range size of Black-backed Woodpeckers (*Picoides arcticus*) in burned ponderosa pine (*Pinus ponderosa*) / Douglas-fir (*Pseudotsuga menziesii*) forests of southwestern Idaho during 2000 and 2002 (6 and 8 years following fire). Home range size for 4 adult males during the post-fledging period was 115.6-420.9 ha using the...

Author(s): Jonathan G. Dudley  
Year Published: 2005  
Type: Document  
Dissertation or Thesis

**Lewis's Woodpecker (*Melanerpes lewis*): a technical conservation assessment**

[www.nrfirescience.org/resource/11498](http://www.nrfirescience.org/resource/11498)

Lewis's woodpecker (*Melanerpes lewis*) is a locally common but patchily distributed woodpecker species usually seen in open forests of western North America. The combination of its sporadic distribution, its diet of adult-stage free-living insects (primarily aerial), its preference to nest in burned landscapes, and its variable...

Author(s): Stephen C. Abele, Victoria A. Saab, Edward O. Garton  
Year Published: 2004  
Type: Document  
Technical Report or White Paper

**Monitoring changes in weed populations: post-fire and post-herbicide treatment**

[www.nrfirescience.org/resource/11040](http://www.nrfirescience.org/resource/11040)

Description not entered

Author(s): Elaine Kennedy Sutherland  
Year Published: 2004  
Type: Document  
Conference Proceedings

**Postfire management on forested public lands of the western United States**

[www.nrfirescience.org/resource/7913](http://www.nrfirescience.org/resource/7913)

Forest ecosystems in the western United States evolved over many millennia in response to disturbances such as wildfires. Land use and management practices have altered these ecosystems, however, including fire regimes in some areas. Forest ecosystems are especially vulnerable to postfire management practices because such practices...

Author(s): Robert L. Beschta, Jonathan J. Rhodes, J. Boone Kauffman, Robert E. Gresswell, G. Wayne Minshall, James R. Karr, David A. Perry, F. Richard Hauer, Christopher A. Frissell  
Year Published: 2004  
Type: Document  
Book or Chapter or Journal Article

**Establishment of aerially seeded big sagebrush following southern Idaho wildfires**

[www.nrfirescience.org/resource/11412](http://www.nrfirescience.org/resource/11412)

In the western United States, big sagebrush (*Artemisia tridentata*) steppe communities dominate approximately 60 million ha (148 million acres) and comprise the largest vegetation type (Wambolt and Hoffman 2001). However, due to the invasion of exotic plants, fire has become a driving force in the ecology and management of sagebrush...

Author(s): Cindy R. Lysne, Michael L. Pellant  
Year Published: 2004  
Type: Document  
Technical Report or White Paper

**Postfire seeding for erosion control: effectiveness and impacts on native plant communities**

[www.nrfirescience.org/resource/7911](http://www.nrfirescience.org/resource/7911)

Large, high-severity wildfires remove vegetation cover and expose mineral soil, often causing erosion

and runoff during postfire rain events to increase dramatically. Land-management agencies in the United States are required to assess site conditions after wildfire and, where necessary, implement emergency watershed rehabilitation...

Author(s): Jan L. Beyers

Year Published: 2004

Type: Document

Book or Chapter or Journal Article, Synthesis

### **Responses of stream benthic macroinvertebrates to fire**

[www.nrfirescience.org/resource/7964](http://www.nrfirescience.org/resource/7964)

Synthesis of published research on the responses of stream benthic macroinvertebrates to fire in western United States indicates a consistent pattern of response that can guide resource management and future research. Direct effects of fire generally are minor or indiscernible. Indirect effects, resulting primarily from increased...

Author(s): G. Wayne Minshall

Year Published: 2003

Type: Document

Book or Chapter or Journal Article

### **On the impact of fire suppression and BAER restoration on weeds**

[www.nrfirescience.org/resource/11043](http://www.nrfirescience.org/resource/11043)

In 2000, wildfires burned more than 200,000 acres on the Bitterroot National Forest of Montana and nearly 1.5 million acres in the Northern and Intermountain Regions. Management activities associated with fire suppression and post-fire restoration have had the unintentional consequence of promoting invasive weeds. As part of fire...

Author(s): Elaine Kennedy Sutherland

Year Published: 2003

Type: Document

Conference Proceedings

### **Quick response small catchment monitoring techniques for comparing postfire rehabilitation treatment effectiveness**

[www.nrfirescience.org/resource/11000](http://www.nrfirescience.org/resource/11000)

Increased runoff and erosion commonly occur after wildfires with the onset of precipitation events. Various erosion mitigation treatments are often used after wildfires to reduce flooding and sedimentation. The effectiveness of these treatments has not been well documented in the literature; therefore we undertook a rapid response...

Author(s): Peter R. Robichaud, Robert E. Brown

Year Published: 2003

Type: Document

Conference Proceedings

### **Selection of fire-created snags at two spatial scales by cavity-nesting birds**

[www.nrfirescience.org/resource/11198](http://www.nrfirescience.org/resource/11198)

We examined the use of snag stands by seven species of cavity-nesting birds from 1994-1998. Selection of snags was studied in logged and unlogged burned forests at two spatial scales: microhabitat (local vegetation characteristics) and landscape (composition and patterning of surrounding vegetation types). We modeled nest occurrence...

Author(s): Victoria A. Saab, Ree Brannon, Jonathan G. Dudley, Larry Donohoo, Dave Vanderzanden, Vicky Johnson, Henry Lachowski

Year Published: 2002

Type: Document  
Technical Report or White Paper

### **Effects of wildfire and post-fire salvage logging on avian communities in conifer-dominated forests of the western United States**

[www.nrfirescience.org/resource/7956](http://www.nrfirescience.org/resource/7956)

Description not entered

Author(s): Natasha B. Kotliar, Sallie Hejl, Richard L. Hutton, Victoria A. Saab, C. P. Melcher, Mary E. McFadzen

Year Published: 2002

Type: Document

Book or Chapter or Journal Article

### **Effects of fire and post-fire salvage logging on avian communities in conifer-dominated forests of the western United States**

[www.nrfirescience.org/resource/14594](http://www.nrfirescience.org/resource/14594)

A one-day symposium organized by the editors was held the following year in Portland, Oregon, at the annual meeting of the Cooper Ornithological Society. The central focus of the symposium was to contrast patterns in the western versus eastern United States, and to differentiate and contrast natural...

Author(s): Natasha B. Kotliar, Sallie Hejl, Richard L. Hutto, Victoria A. Saab, C. P. Melcher, Mary E. McFadzen

Year Published: 2002

Type: Document

Conference Proceedings

### **Fire and invasive species within the temperate and boreal coniferous forests of western North America**

[www.nrfirescience.org/resource/10966](http://www.nrfirescience.org/resource/10966)

Invasive, nonnative plant species have been a concern of land managers within the temperate and boreal coniferous forest eco-region for nearly a century. Fire management, timber harvest, grazing, mining, recreation, and agriculture have not only exacerbated invasive species establishment and spread, but have been impacted by such...

Author(s): Richy J. Harrod, Sarah Reichard

Year Published: 2000

Type: Document

Conference Proceedings, Synthesis

### **Evaluating the effectiveness of postfire rehabilitation treatments**

[www.nrfirescience.org/resource/11194](http://www.nrfirescience.org/resource/11194)

Spending on postfire emergency watershed rehabilitation has increased during the past decade. A west-wide evaluation of USDA Forest Service burned area emergency rehabilitation (BAER) treatment effectiveness was undertaken as a joint project by USDA Forest Service Research and National Forest System staffs. This evaluation covers...

Author(s): Peter R. Robichaud, Jan L. Beyers, Daniel G. Neary

Year Published: 2000

Type: Document

Technical Report or White Paper

### **Responses of cavity-nesting birds to stand-replacement fire and salvage logging in ponderosa pine/douglas-fir forests of southwestern Idaho**

[www.nrfirescience.org/resource/11413](http://www.nrfirescience.org/resource/11413)

From 1994 to 1996, researchers monitored 695 nests of nine cavity-nesting bird species and measured vegetation at nest sites and at 90 randomly located sites in burned ponderosa pine forests of southwestern Idaho. Site treatments included two types of salvage logging, and unlogged controls. All bird species selected nest sites with...

Author(s): Victoria A. Saab, Jonathan G. Dudley

Year Published: 1998

Type: Document

Technical Report or White Paper

### **Length and timing of grazing on postburn productivity of two bunchgrasses in an Idaho experimental range**

[www.nrfirescience.org/resource/8213](http://www.nrfirescience.org/resource/8213)

Plant mortality and productivity in semiarid grasslands may be affected by the length of time grazing is excluded during the postfire regeneration period. The degree of grazing tolerance for the semiarid bunchgrass species, *Festuca idahoensis* and *Agropyron spicatum*, exposed to fire, and how the variation in grazing tolerance was...

Author(s): Stephen C. Bunting, Ronald Robberecht, Guillermo E. Defosse

Year Published: 1998

Type: Document

Book or Chapter or Journal Article

### **Vegetal recovery following wildfire in seeded and unseeded sagebrush steppe**

[www.nrfirescience.org/resource/11459](http://www.nrfirescience.org/resource/11459)

Following an August wildfire, sagebrush (*Artemisia L.*)/grass benchlands adjacent to Pocatello, Ida., were seeded with a mixture of exotic wheatgrasses and forbs by rangeland drill in November 1987. The effects of seeding on vegetation development in the immediate postfire years were evaluated by comparing plant density, vegetal...

Author(s): Teresa D. Ratzlaff, Jay E. Anderson

Year Published: 1995

Type: Document

Book or Chapter or Journal Article

### **Deterioration of fire-killed and fire-damaged timber in the Western United States**

[www.nrfirescience.org/resource/11159](http://www.nrfirescience.org/resource/11159)

Fire-killed and fire-damaged timber are an important source of fiber and are becoming more important because of a decrease in the land base available for timber harvest. Forest managers need to know the causes of deterioration and degrade, the expected losses in product volume and value, and the impact of time on deterioration. This...

Author(s): Eini C. Lowell, Susan A. Willits, Robert L. Krahmer

Year Published: 1992

Type: Document

Technical Report or White Paper

### **Fire's effects on a small bird population**

[www.nrfirescience.org/resource/11188](http://www.nrfirescience.org/resource/11188)

Changes in bird populations as a result of a 122 ha forest fire are evaluated. There is little evidence of any drastic effect on numbers of birds, species, or species diversity in the year of the fire or 2 years later.

Author(s): L. Jack Lyon, John M. Marzluff

Year Published: 1985

Type: Document  
Technical Report or White Paper

### **Early postfire revegetation in a western Montana Douglas-fir forest**

[www.nrfirescience.org/resource/11960](http://www.nrfirescience.org/resource/11960)

Development of natural vegetation and seeded grasses on a severely burned Douglas-fir forest area is described for the first 5 postfire years. Results are described separately for ravine and upland sites.

Results of special studies of moss recovery and tree seedling distribution are also reported.

Author(s): Marilyn F. Crane, James R. Habeck, William C. Fischer

Year Published: 1984

Type: Document

Technical Report or White Paper

### **The Sleeping Child Burn - 21 years of postfire change**

[www.nrfirescience.org/resource/11961](http://www.nrfirescience.org/resource/11961)

In early August 1961, more than 26,000 acres (10,500 ha) of upper montane and subalpine forest on the Bitterroot National Forest burned in a lightning-caused wildfire. At the time, the Sleeping Child Burn represented the single largest forest fire in the Northern Rocky Mountains in more than 20 years.

Historically, large wildfires...

Author(s): L. Jack Lyon

Year Published: 1984

Type: Document

Technical Report or White Paper

### **Vegetal development on the Sleeping Child burn in western Montana, 1961 to 1973**

[www.nrfirescience.org/resource/11951](http://www.nrfirescience.org/resource/11951)

In the year following the 1961 Sleeping Child forest fire on the Bitterroot National Forest, Montana, 11 permanent transects were established within the burn. Vegetation development was recorded through 1973, but only four transects were considered indicative of seral forest succession independent of superimposed management...

Author(s): L. Jack Lyon

Year Published: 1976

Type: Document

Technical Report or White Paper

### **Hayman fire: short- and long-term geomorphic change and recovery**

[www.nrfirescience.org/resource/13027](http://www.nrfirescience.org/resource/13027)

Lee MacDonald, Professor, Colorado State University, Department of Forest, Rangeland, and Watershed Stewardship, discusses geomorphic changes following the Hayman and Schoonover wildfires at the Hayman Fire Science Symposium: Lessons Learned after Ten Years of Recovery, Rehabilitation, and Restoration.

Type: Media

Webinar

### **Post-fire tree mortality and management**

[www.nrfirescience.org/resource/14214](http://www.nrfirescience.org/resource/14214)

This presentation was recorded during the 2016 State of the State and Forest Health Conference in Corvallis, OR.

Type: Media



Video

**Linking basic and applied research, multi-resource management, public education, and enforcement: post-fire archeology on the Shoshone National Forest**

[www.nrfirescience.org/resource/13738](http://www.nrfirescience.org/resource/13738)

Especially in remote, Wilderness settings, fires produce a complex array of both direct and indirect impacts to heritage resources that creates a cascade of complex research and management issues and opportunities. Over the last decade we have been working to align goals of academic research programs and...

Type: Media

Video

**Burned area recovery project: a stakeholder perspective**

[www.nrfirescience.org/resource/13320](http://www.nrfirescience.org/resource/13320)

In this video, Larry Campbell, member of Friends of the Bitterroot, spoke about his organization's opposition to the salvage logging component of the BNF's Burned Area Recovery project following the Fires of 2000. This was filmed along Rye Creek, which was one of the stops during the Fires of 2000 field trip that was part of...

Type: Media

Video

**Burned area emergency response**

[www.nrfirescience.org/resource/13309](http://www.nrfirescience.org/resource/13309)

In this video, Marilyn Wildey, Hydrology Technician with the Bitterroot National Forest, describes the Burned Area Emergency Response following the Bitterroot fires of 2000. This was filmed at the Bitterroot National Forest headquarters in Hamilton, MT, which was one of the stops during the Fires of 2000 field trip that was part of...

Type: Media

Video

**Post-fire reforestation considerations**

[www.nrfirescience.org/resource/13221](http://www.nrfirescience.org/resource/13221)

Post-fire environments are dynamic and complex and trends of recent fires within the Sierra mixed conifer type include uncharacteristically large areas of high vegetation burn severity. These trends add to the complexity of opportunities and challenges for post-fire restoration efforts - a key component of which is reforestation....

Type: Media

Webinar

**Post-wildfire seeding in forests of the West: effectiveness, trends over time, and fire management perspectives**

[www.nrfirescience.org/resource/13037](http://www.nrfirescience.org/resource/13037)

Dr. Pete Fule presented results from the Joint Fire Science Program (JFSP) project synthesizing existing information on post-wildfire seeding. The webinar covered key findings from an evidence-based systematic review conducted to examine the effectiveness and effects of post-fire seeding treatments on soil stabilization and plant...

Type: Media

Webinar

**10 years of post-fire treatment monitoring - Learning about soil and vegetation recovery**

[www.nrfirescience.org/resource/12937](http://www.nrfirescience.org/resource/12937)

Following the 2005 School Fire that burned about 50,000 acres of forests and grasslands on the Umatilla National Forest, Washington, managers wanted to limit weed spread and soil erosion in severely burned areas. Various mulch treatments (wheat straw, wood strand, and hydromulch) were used to control erosion on steep slopes above...

Type: Media

Webinar

### **Who's to blame? Fire management in mixed-ownership landscapes**

[www.nrfirescience.org/resource/15097](http://www.nrfirescience.org/resource/15097)

Fuels are the only component of the fire triangle that forest and fire managers can alter to change fire behavior. There have been numerous studies examining how fuel reduction treatments and salvage logging alter fire behavior, severity, and its' ecological impacts. However, less attention has been paid to how different forest...

Type: Media

Webinar

### **Post-fire restoration considerations**

[www.nrfirescience.org/resource/14074](http://www.nrfirescience.org/resource/14074)

Post-fire environments are dynamic and complex and trends of recent fires within the Sierra mixed conifer type include uncharacteristically large areas of high vegetation burn severity. These trends add to the complexity of opportunities and challenges for post-fire restoration efforts - a key component of which is...

Type: Media

Webinar

### **Quaking aspen: a burning desire in an 'asbestos forest'**

[www.nrfirescience.org/resource/13673](http://www.nrfirescience.org/resource/13673)

Speaker: Paul Rogers, Director, Western Aspen Alliance, Adjunct Associate Professor, Utah State University. Event: Restoring the West Conference 2015 - Restoration and Fire in the Interior West.

Type: Media

Video

### **Burned area recovery project: the Bitterroot National Forest experience**

[www.nrfirescience.org/resource/13312](http://www.nrfirescience.org/resource/13312)

In this video, Marilyn Wildey, Hydrology Technician with the Bitterroot National Forest, describes the Burned Area Recovery Project proposed after the Bitterroot fires of 2000. This was filmed at the Bitterroot National Forest headquarters in Hamilton, MT, which was one of the stops during the Fires of 2000 field trip that was part...

Type: Media

Video

### **Ten years of post-fire treatment monitoring - Learning about soil and vegetation recovery**

[www.nrfirescience.org/resource/13234](http://www.nrfirescience.org/resource/13234)

Following the 2005 School Fire that burned about 50,000 acres of forests and grasslands on the Umatilla National Forest, Washington, managers wanted to limit weed spread and soil erosion in severely burned areas. Various mulch treatments (wheat straw, wood strand, and hydromulch) were used to control erosion on steep slopes above...

Type: Media

Webinar

**Climate, weather, and sagebrush seed sources: experimental insights on challenges and opportunities**

[www.nrfirescience.org/resource/13219](http://www.nrfirescience.org/resource/13219)

Matt Germino, Research Ecologist, USGS Snake River Field Station, Boise, ID, discusses experimental insights on challenges and opportunities regarding climate, weather, and sagebrush seed sources.

Type: Media

Webinar